

## Numerical Analysis - Approximating Functions

*Latest Modification: August 27, 2001*

**Notation:** Equation numbers are placed in parentheses, (**Eq. 1**), at the end of the sentence preceding the mathematical presentation of the equation. They should not be read as if they are part of the sentence. When reference is made to an equation, the word "equation" will be spelled out and not abbreviated. Mathematical symbols are not in all cases precise. Greek symbols will show as such on a Windows operating system, but not a Unix or Mac system.

### Series Approximation

Given values of a function  $f(x)$  at a finite number of discrete points ( $n$ ),

$$x_0, \quad x_1, \quad x_2, \quad x_3, \quad \dots, \quad x_{n-1}, \quad x_n$$

$$f(x_0), f(x_1), f(x_2), f(x_3), \dots, f(x_{n-1}), f(x_n)$$

the goal is to construct an arbitrary, but simpler, function  $p(x)$  in the form of series that takes on the values of  $f_i$  at the  $n$  points  $x_i$ , that is (**Eq. 1**),

$$p(x) \cong f(x), \text{ for } x_0 < x < x_n$$

and possibly for  $x < x_0$  or  $x > x_n$

that is to say (**Eq. 2**)

$$p(x) = f(x_i), \text{ for } x_0, \dots, x_n$$

The function  $p(x)$  can take a variety of forms, such as

- when  $p(x)$  is a polynomial, the process of representing  $f(x)$  by  $p(x)$  is called a **polynomial approximation**,
- when  $p(x)$  is a finite trigonometric series, the process of representing  $f(x)$  by  $p(x)$  is called a **trigonometric approximation**, and
- in like manner  $p(x)$  may be a series of:
  - exponential functions
  - Legendre polynomials
  - Bessel functions
  - etc.

Approximating functions are the basis for interpolation schemes, and interpolation schemes are the basis for numerical quadrature relations. Thus approximating functions represent an important element in numerical methods.

## Difference Tables

**B**efore we consider polynomial interpolation schemes, we should develop the concept of difference tables in the following document called [Difference Tables](#).

---

## Interpolation

**N**ow that we have the concept of differences and their notation, we can develop the most elementary interpolation schemes known as polynomial interpolation found in the following document [Polynomial Interpolation](#).

**A** frequently encounter interpolation problem is one in which the data is not equally spaced in the independent variable. This leads us to consider the material in the following document on [Non-Equidistant Interpolation](#).

---

## Least Square Data Analysis

**F**inally, we should explore a different approach to approximating functions known as least squares as discussed in the following document [Least Squares](#).

---

*Physics & Astronomy Department, George Mason University  
Maintained by J. C. Evans; [jevans@gmu.edu](mailto:jevans@gmu.edu)*

**PHYS 251 - Introduction to Computer Techniques in Physics****Difference Tables***Latest Modification: September 28, 1999*

**Notation:** Equation numbers are placed in parentheses, (**Eq. 1**), at the end of the sentence preceding the mathematical presentation of the equation. They should not be read as if they are part of the sentence. When reference is made to an equation, the word "equation" will be spelled out and not abbreviated. Mathematical symbols are not in all cases precise. Greek symbols will show as such on a Windows operating system, but not a Unix or Mac system.

**Horizontal Difference Table**

Difference tables lend themselves to the use of spreadsheets as a means of solving many problems rather than resorting to writing computer code in either Fortran or C/C++. One type of difference table is known as a **horizontal difference table** in which differences are maintained on horizontal rows with the function as shown below for which successive differences through the fourth are denoted by  $\Delta_1 f_i(x)$ ,  $\Delta_2 f_i(x)$ ,  $\Delta_3 f_i(x)$ , and  $\Delta_4 f_i(x)$ .

$x_i$	$f_i(x)$	$\Delta_1 f_i(x)$	$\Delta_2 f_i(x)$	$\Delta_3 f_i(x)$	$\Delta_4 f_i(x)$
...	...	...	...	...	...
$x_{-2}$	$f_{-2}$	$f_{-2} - f_{-3}$	$f_{-2} - 2f_{-3} + f_{-4}$	...	...
$x_{-1}$	$f_{-1}$	$f_{-1} - f_{-2}$	$f_{-1} - 2f_{-2} + f_{-3}$	$f_{-1} - 3f_{-2} + 3f_{-3} - f_{-4}$	...
$x_0$	$f_0$	$f_0 - f_{-1}$	$f_0 - 2f_{-1} + f_{-2}$	$f_0 - 3f_{-1} + 3f_{-2} - f_{-3}$	$f_0 - 4f_{-1} + 6f_{-2} - 4f_{-3} + f_{-4}$
$x_1$	$f_1$	$f_1 - f_0$	$f_1 - 2f_0 + f_{-1}$	$f_1 - 3f_0 + 3f_{-1} - f_{-2}$	$f_1 - 4f_0 + 6f_{-1} - 4f_{-2} + f_{-3}$
$x_2$	$f_2$	$f_2 - f_1$	$f_2 - 2f_1 + f_0$	$f_2 - 3f_1 + 3f_0 - f_{-1}$	$f_2 - 4f_1 + 6f_0 - 4f_{-1} + f_{-2}$
$x_3$	$f_3$	$f_3 - f_2$	$f_3 - 2f_2 + f_1$	$f_3 - 3f_2 + 3f_1 - f_0$	$f_3 - 4f_2 + 6f_1 - 4f_0 + f_{-1}$
$x_4$	$f_4$	$f_4 - f_3$	$f_4 - 2f_3 + f_2$	$f_4 - 3f_3 + 3f_2 - f_1$	$f_4 - 4f_3 + 6f_2 - 4f_1 + f_0$
...	...	...	...	...	...

## Error Propagation in Difference Tables

Errors propagate in a horizontal difference table as shown below, where  $\epsilon$  is a small increment in the function  $f$ .

$x_i$	$f_i(x)$	$\Delta_1 f_i(x)$	$\Delta_2 f_i(x)$	$\Delta_3 f_i(x)$	$\Delta_4 f_i(x)$
...	...	...	...	...	...
$x_{-2}$	$f_{-2}$	...	...	...	...
$x_{-1}$	$f_{-1}$	...	...	...	...
$x_0$	$f_0(x) + \epsilon$	$\Delta_1 f_0(x) + \epsilon$	$\Delta_2 f_0(x) + \epsilon$	$\Delta_3 f_0(x) + \epsilon$	$\Delta_4 f_0(x) + \epsilon$
$x_1$	$f_1$	$\Delta_1 f_1(x) - \epsilon$	$\Delta_2 f_1(x) - 2\epsilon$	$\Delta_3 f_1(x) - 3\epsilon$	$\Delta_4 f_1(x) - 4\epsilon$
$x_2$	$f_2$	...	$\Delta_2 f_2(x) + \epsilon$	$\Delta_3 f_2(x) + 3\epsilon$	$\Delta_4 f_2(x) + 6\epsilon$
$x_3$	$f_3$	...	...	$\Delta_3 f_3(x) - \epsilon$	$\Delta_4 f_3(x) - 4\epsilon$
$x_4$	$f_4$	...	...	...	$\Delta_4 f_4(x) + \epsilon$
...	...	...	...	...	...

There are several points one should note about the propagation of the error.

1. The error moves to successive rows with successive differences.
2. The coefficients of  $\epsilon$ 's are binomial coefficients with alternating signs.
3. The algebraic sum of the  $\epsilon$ 's in any difference column is zero.
4. The first erroneous difference of any order is on the same horizontal line as the erroneous functional value.

## Polynomial Interpolation

---

*Latest Modification: November 1, 2001*

---

**Notation:** Equation numbers are placed in parentheses, (**Eq. 1**), at the end of the sentence preceding the mathematical presentation of the equation. They should not be read as if they are part of the sentence. When reference is made to an equation, the word "equation" will be spelled out and not abbreviated. Mathematical symbols are not in all cases precise. Greek symbols will show as such on a Windows operating system, but not a Unix or Mac system.

---

### Polynomial Interpolation

The justification for approximating with polynomials rest on a theorem by Weierstrass in 1885, to which

"...every continuous function in an interval (a,b) can be represented in that interval to any desired accuracy by a polynomial."

One can also show that the following propositions are true for a polynomial

1. nth differences of an nth-degree polynomial are constant for an arithmetic progression (equal interval) of the independent variable,
  2. if nth differences of a tabulated function for equal interval data are constant, the unknown function is a polynomial of degree n.
- 

### Example

Let  $f(x) = x^2 + 3x + 2$ . Forming differences in a horizontal difference table, we have the table shown below.

$x_i$	$f_i(x)$	$\Delta_1 f_i(x)$	$\Delta_2 f_i(x)$	$\Delta_3 f_i(x)$	$\Delta_4 f_i(x)$
0.000000	2.000000				
1.000000	6.000000	4.000000			
2.000000	12.000000	6.000000	2.000000		
3.000000	20.000000	8.000000	2.000000	0.000000	
4.000000	30.000000	10.000000	2.000000	0.000000	
5.000000	42.000000	12.000000	2.000000	0.000000	
6.000000	56.000000	14.000000	2.000000	0.000000	

Since the function is indeed a 2nd degree polynomial the 2nd differences are constant.

## Polynomial Interpolation Retaining Second Differences

If we restrict our consideration to the retention of second differences and ignore all higher-order differences, there are four useful polynomial approximations that can be derived for equal-interval interpolation.

**Newton's Forward Interpolation:** Given values of a function  $f(x)$  at a finite number of discrete points ( $n$ ),

$$x_0, \quad x_1, \quad x_2, \quad x_3, \quad \dots, \quad x_{n-1}, \quad x_n$$

$$f(x_0) = f_0, \quad f(x_1) = f_1, \quad f(x_2) = f_2, \quad f(x_3) = f_3, \quad \dots, \quad f(x_{n-1}) = f_{n-1}, \quad f(x_n) = f_n$$

and letting  $u = [(x - x_i) / (x_{i+1} - x_i)]$ , then Newton's forward interpolation formula can be expressed using difference notation as

$$p(x) = p(x_0 + u\Delta x)$$

or substituting the difference values of the function, we have

$$p(x) = f_0 + u\Delta_1 f_1 + [u(u-1)/2!]\Delta_2 f_2 + [u(u-1)(u-2)/3!]\Delta_3 f_3 + \dots + [u(u-1)(u-2)\dots(u-n+1)/n!]\Delta_n f_n + \dots$$

which can be simplified by retaining only second order differences and substituting the functional values for the differences (**Eq. 1**)

$$p(x) = f_0 + u(f_1 - f_0) + [u(u-1)/2](f_2 - 2f_1 + f_0)$$

Since Newton's forward interpolation formula uses values of the function at points ahead in

the table, it can be used at the beginning of the tabulated values, but not at the end.

**Newton's Backward Interpolation:** Given values of a function  $f(x)$  at a finite number of discrete points ( $n$ ), and letting  $u = [(x - x_i) / (x_i - x_{i-1})]$ , then Newton's backward interpolation formula can be expressed using difference notation as

$$p(x) = p(x_0 + u\Delta x)$$

or substituting the difference values of the function, we have

$$p(x) = f_n + u\Delta_1 f_n + [u(u+1)/2!]\Delta_2 f_n + [u(u+1)(u+2)/3!]\Delta_3 f_n + \dots + [u(u+1)(u+2)\dots(u+n-1)/n!]\Delta_n f_n + \dots$$

which can be simplified by retaining only second order differences and substituting the functional values for the differences (**Eq. 2**)

$$p(x) = f_0 + u(f_0 - f_{-1}) + [u(u+1)/2](f_0 - 2f_{-1} + f_{-2})$$

which is useful for interpolating near the end of the tabulated functions.

**Stirling's Central-Difference Interpolation:** This interpolation formula is based on a diagonal difference table rather than a horizontal difference table. Without going in to the distinctions between the two types of difference tables, we can write down an interpolation formula including only second differences that is similar to Newton's interpolation formulas. Thus given values of a function  $f(x)$  at a finite number of discrete points ( $n$ ), and letting  $u = [(x - x_i) / (x_{i+1} - x_i)]$ , Stirling's central-difference interpolation formula can be expressed using functional values as (**Eq. 3**)

$$p(x) = f_0 + (u/2)(f_1 - f_{-1}) + [u^2/2](f_1 - 2f_0 + f_{-1})$$

which also uses three successive points to approximate the function as does Newton.

**Bessel's Central-Difference Interpolation:** This interpolation formula is also based on a diagonal difference table rather than a horizontal difference table. We shall also write down an interpolation formula including only second differences that is similar to Newton's and Stirling's interpolation formulas. Thus given values of a function  $f(x)$  at a finite number of discrete points ( $n$ ), and letting  $u = [(x - x_i) / (x_{i+1} - x_i)]$ , Bessel's central-difference interpolation formula can be expressed using functional values as (**Eq. 4**)

$$p(x) = f_0 + u(f_1 - f_0) + [u(u-1)/4](f_2 - f_1 - f_0 + f_{-1})$$

which uses four successive points rather than three to approximate the function. Note that in equations 1, 2, 3, and 4, there is a zero-order term, a first-order term, and a second-order term. The zero- and first-order terms constitute essentially a linear interpolation, while the second-order term is a correction term to that linear interpolation.

## Example

Find  $f(x) = \exp(-x)$ , at the point  $x = 1.7565$ , given the following values of the function.

$i$	$x_i$	$f_i$
-2	1.730000000	0.177284100
-1	1.740000000	0.175520401
0	1.750000000	0.173773944
1	1.760000000	0.172044864
2	1.770000000	0.170332989

Forming the value of  $u$  for the four equidistance interpolation formulas, we have for Newton's forward, Stirling's, and Bessel's interpolation formulas

$$u = (x - x_0) / (x_1 - x_0) = (1.7565 - 1.75) / (1.76 - 1.75) = 0.0065 / 0.01 = +0.65,$$

**Newton's Forward Interpolation:**

$$u(u-1)/2 = 0.65(0.65-1)/2 = -0.11375,$$

**Stirling's Central-Difference Interpolation:**

$$u/2 = 0.65/2 = 0.325,$$

$$u^2/2 = (0.65)^2/2 = 0.21125,$$

**Bessel's Central-Difference Interpolation:**

$$u(u-1)/4 = 0.65(0.65-1)/4 = -0.056875.$$

For Newton's backward formula, the value of  $u$  is given by

$$u = (x - x_1) / (x_1 - x_0) = (1.7565 - 1.76) / (1.76 - 1.75) = -0.0035 / 0.01 = -0.35,$$

**Newton's Backward Interpolation:**

$$u(u+1)/2 = -0.35(-0.35+1)/2 = -0.11375.$$

The following table evaluates the three terms that appear in each of the four interpolation formula. Note that in order to apply Newton's backward formula, we must re-label the points so that  $i = -1, 0, 1$  become  $i = -2, -1, 0$ .



Interpolation Formula	Zero-Order Term	First-Order Term	Second-Order Term	f(u)
Newton's Forward	0.173773944	-0.001123902	-0.000001957	0.172648085
Newton's Backward	0.172044864	+0.000605178	-0.000001977	0.172648065
Stirling's Formula	0.173773944	-0.001129550	+0.000003671	0.172648065
Bessel's Formula	0.173773944	-0.001123902	-0.000001967	0.172648075
Exact Solution				0.172648076

It is evident that Bessel's interpolation formula has given us the most accurate result. This is in part true because Bessel's formula is known to be more accurate when  $u$  lies between 0.25 and 0.75.

### Non-Equidistant Interpolation

A frequently encountered interpolation problem is one in which the data is not equally spaced in the independent variable. This leads us to consider Lagrangian interpolation in the following document on [Non-Equidistant Interpolation](#).

*Physics & Astronomy Department, George Mason University  
Maintained by J. C. Evans; [jevans@gmu.edu](mailto:jevans@gmu.edu)*

# Non-Equidistant Interpolation

## Lagrange's Formula

---

*Latest Modification: November 1, 2001*

---

**Notation:** Equation numbers are placed in parentheses, (**Eq. 1**), at the end of the sentence preceding the mathematical presentation of the equation. They should not be read as if they are part of the sentence. When reference is made to an equation, the word "equation" will be spelled out and not abbreviated. Mathematical symbols are not in all cases precise. Greek symbols will show as such on a Windows operating system, but not a Unix or Mac system.

---

### Non-Equidistant Data Interpolation

The previous discussion of interpolation is predicated on equidistant intervals in the independent variable  $x$ . Sometimes it is inconvenient or impossible to obtain function values on an equidistant interval. In such cases, Lagrange's formula can be fashioned to only utilize such data as may be available.

The general form of Lagrange's interpolation formula is given by (**Eq. 1**)

$$p(x) = \sum_{j=0}^n \left\{ \left[ \prod_{i=0, i \neq j}^n (x - x_i) \right] / \left[ \prod_{i=0, i \neq j}^n (x_j - x_i) \right] \right\} f_j$$

where

$$\prod_{i=0}^n (x - x_i) = (x - x_1)(x - x_2) \dots (x - x_n)$$

$$\prod_{i=0, i \neq j}^n (x_j - x_i) = (x_j - x_0) \dots (x_j - x_{i-1})(x_j - x_{i+1}) \dots (x_j - x_n)$$

We should note that

1. Lagrange's formula does not involve successive differences as does Newton's, Stirling's and Bessel's interpolation formulas.
  2. Lagrange's formula does involve  $n+1$  successive pairs of variables  $(x_n, f_n)$ .
  3. Although Lagrange's formula works for non-equidistant data, it is not restricted to such a limitation, i.e., it can be applied to equidistant data.
  4. If we know rates of change (differences) or rates of rates of change of the function, their neglect is to ignore information.
- 

### Cubic Lagrangian Interpolation

Suppose one wishes to construct a cubic polynomial through four successive points of a

data set which we will label as  $(x_1, f_1)$ ,  $(x_2, f_2)$ ,  $(x_3, f_3)$ , and  $(x_4, f_4)$ . This can be accomplished from our general relation for  $n+1$  points where we can construct the following relation (Eq. 2)

$$p(x) = l_{1,4} f_1 + l_{2,4} f_2 + l_{3,4} f_3 + l_{4,4} f_4$$

where

$$l_{1,4} = [(x - x_2)(x - x_3)(x - x_4)] / [(x_1 - x_2)(x_1 - x_3)(x_1 - x_4)]$$

$$l_{2,4} = [(x - x_1)(x - x_3)(x - x_4)] / [(x_2 - x_1)(x_2 - x_3)(x_2 - x_4)]$$

$$l_{3,4} = [(x - x_1)(x - x_2)(x - x_4)] / [(x_3 - x_1)(x_3 - x_2)(x_3 - x_4)]$$

$$l_{4,4} = [(x - x_1)(x - x_2)(x - x_3)] / [(x_4 - x_1)(x_4 - x_2)(x_4 - x_3)]$$

If one wishes, this expression can be written in the form (with a little algebra) as

$$p(x) = ax^3 + bx^2 + cx + d$$

However, from the computational point of view there is no real advantage in doing so.

## Example

Find the cubic polynomial whose graph contains the four successive points (0,1), (1,2), (2,0), and (3,-2). Setting  $x_1 = 0$ ,  $f_1 = 1$ ,  $x_2 = 1$ ,  $f_2 = 2$ ,  $x_3 = 2$ ,  $f_3 = 0$ , and  $x_4 = 3$ ,  $f_4 = -2$ , we can form the values of the  $l$ s

$$\begin{aligned} l_{1,4} &= [(x - x_2)(x - x_3)(x - x_4)] / [(x_1 - x_2)(x_1 - x_3)(x_1 - x_4)] \\ &= [(x - 1)(x - 2)(x - 3)] / [(0 - 1)(0 - 2)(0 - 3)] \\ &= [(x - 1)(x - 2)(x - 3)] / -6 \end{aligned}$$

$$\begin{aligned} l_{2,4} &= [(x - x_1)(x - x_3)(x - x_4)] / [(x_2 - x_1)(x_2 - x_3)(x_2 - x_4)] \\ &= [(x - 0)(x - 2)(x - 3)] / [(1 - 0)(1 - 2)(1 - 3)] \\ &= [x(x - 2)(x - 3)] / 2 \end{aligned}$$

$$\begin{aligned} l_{3,4} &= [(x - x_1)(x - x_2)(x - x_4)] / [(x_3 - x_1)(x_3 - x_2)(x_3 - x_4)] \\ &= [(x - 0)(x - 1)(x - 3)] / [(2 - 0)(2 - 1)(2 - 3)] \\ &= [x(x - 1)(x - 3)] / -2 \end{aligned}$$

$$\begin{aligned} l_{4,4} &= [(x - x_1)(x - x_2)(x - x_3)] / [(x_4 - x_1)(x_4 - x_2)(x_4 - x_3)] \\ &= [(x - 0)(x - 1)(x - 2)] / [(3 - 0)(3 - 1)(3 - 2)] \\ &= [x(x - 1)(x - 2)] / 6. \end{aligned}$$

Inserting the values for the  $l$ s into Equation 2, we have

$$\begin{aligned} f(x) &= l_{1,4} f_1 + l_{2,4} f_2 + l_{3,4} f_3 + l_{4,4} f_4 \\ &= \{ [(x - 1)(x - 2)(x - 3)] / -6 \} 1 + \{ [x(x - 2)(x - 3)] / 2 \} 2 + \{ [x(x - 1)(x - 3)] / -2 \} 0 + \{ [x(x - 1)(x - 2)] / 6 \} (-2) \end{aligned}$$

$$= (-1/6) [(x-1)(x-2)(x-3)] + [x(x-2)(x-3)] + (-1/3) [x(x-1)(x-2)] .$$

Multiplying out the terms and collecting yields the desired polynomial

$$f(x) = 1/2 x^3 - 3x^2 + 7/2 x + 1 .$$

---

*Physics & Astronomy Department, George Mason University*  
*Maintained by J. C. Evans; jevans@gmu.edu*

## Least Squares

*Latest Modification: November 14, 2001*

**Notation:** Equation numbers are placed in parentheses, (**Eq. 1**), at the end of the sentence preceding the mathematical presentation of the equation. They should not be read as if they are part of the sentence. When reference is made to an equation, the word "equation" will be spelled out and not abbreviated. Mathematical symbols are not in all cases precise. Greek symbols will show as such on a Windows operating system, but not a Unix or Mac system.

### Defining a "Close Fitting" Approximating Function

Earlier we had discussed approximating functions ([Approximating Functions](#)). In that discussion the approximating function contained all of the points of a given discrete function, i.e., the function was constructed such that it agreed precisely with the discrete points used in its construction. Suppose now we wish to construct approximating functions whose graphs need not necessarily contain the points of given discrete function, but which "fits the function closely." Such is the method known as least squares.

In order to understand this idea of "fits the function closely," let us consider a simple example. Given the points ( 0, -1 ), ( 1, 1 ), ( 2, 3 ), and ( 3, 4 ), let us find the straight line  $f(x) = a_1 + a_2 x$ , which in some sense "fits the data closely."


Figure 1 shows a graph of the four points that we wish to fit. One means of determining what is a close fit, is to minimize the vertical distances between the given points and the constructed line. Let us calculate the "errors" or "how far the line misses the given points" as follows (**Eq. 1**)

$$\epsilon_1 = a_1 + a_2 x_1 - f_1,$$

$$\epsilon_2 = a_1 + a_2 x_2 - f_2,$$

$$\epsilon_3 = a_1 + a_2 x_3 - f_3,$$

$$\epsilon_4 = a_1 + a_2 x_4 - f_4.$$



where our object is to minimize the total error. We could do that by considering the total error  $E$ . Let us first consider just the sum of the individual errors as given by (**Eq. 2**)

However, a simple sum will not suffice as the following example illustrates. Suppose the values of the errors are as shown

$$E = (0.1) + (-0.1) + (0.2) + (-0.2) = 0 .$$

Although the sum gives us a minimum, it is meaningless since clearly the total error is not zero. Another possible means of minimizing is to take the absolute values of the errors, but this means that the function  $E$  is not always differentiable. To preserve the differentiability of  $E$  and yet avoid the sign problem, we can minimize the squares of the errors as in **(Eq. 3)**

$$E = \epsilon_1^2 + \epsilon_2^2 + \epsilon_3^2 + \epsilon_4^2 .$$

Equation 3 is the desirable definition of  $E$  since (i) it is always differentiable, (ii) individual terms are non-negative, and (iii) because of the parabolic structure  $E$  always has a unique minimum.

---

## Minimizing the Total Error

Let us construct our total error function for the given example **(Eq. 4)**

$$E = [a_1 + 1]^2 + [(a_1 + a_2) - 1]^2 + [(a_1 + 2a_2) - 3]^2 + [(a_1 + 3a_2) - 4]^2 .$$

In order to minimize, we want to set the partial derivative of  $E$  with respect to each of the parameters in the linear fit,  $a_1$  and  $a_2$ , as shown **(Eq. 5)**

$$\delta E / \delta a_1 = 0 , \delta E / \delta a_2 = 0 .$$

which yields the two linear algebraic equations

$$\begin{aligned} 4 a_1 + 6 a_2 &= 7 , \\ 6 a_1 + 14 a_2 &= 19 , \end{aligned}$$

whose solution is

$$\begin{aligned} a_1 &= -0.8 , \\ a_2 &= 1.7 . \end{aligned}$$

Thus the approximating function that "fits the data closely" is (Eq. 6)

$$f(x) = a_1 + a_2 x = -0.8 + 1.7x,$$

and the sense in which it fits closely is that it minimizes the squares of the vertical distances between the approximating function and the discrete points.




Figure 2 shows a graph of the four points that we wish to fit along with approximating function that minimizes the squares of the vertical distances between the given points and the constructed line.

Are there other ways to define "fits closely?" Yes, for example, minimize the squares of the perpendicular distances rather than the vertical distances. However, this is a considerably harder problem to solve. Therefore, we will restrict ourselves to the standard definition of the least squares technique and minimize the vertical distances.

## Periodic or Quasi-Periodic Approximating Functions in Least Squares.

Let us consider a second example, which reveals more of the subtle aspects of the method of least squares.

the general description of the method of least squares is as follows. Given the set of discrete points  $(x_1, f_1)$ ,  $(x_2, f_2)$ ,  $(x_3, f_3)$ , ...  $(x_n, f_n)$ , let  $g(x)$  be a continuous function chosen such as to minimize the squares of the vertical distances, that is, minimize the total error  $E$  (Eq. 7)

$$E = [g(x_1) - f_1]^2 + [g(x_2) - f_2]^2 + [g(x_3) - f_3]^2 + \dots + [g(x_n) - f_n]^2.$$

assuming that  $g(x)$  contains  $m$  independent parameters  $a_1, a_2, \dots, a_m$ . In order to minimize  $E$  with respect to each of the  $m$  independent parameters, we solve the system of algebraic equations given by (Eq. 8)

$$\delta E / \delta a_1 = 0, \delta E / \delta a_2 = 0, \delta E / \delta a_3 = 0, \dots, \delta E / \delta a_m = 0.$$


to yield the desired values of the  $m$  independent parameters  $a_1, a_2, \dots, a_m$ .

Now let us consider the oscilloscope trace shown in Figure 3. For our oscilloscope trace, what is the functional form for  $g(x)$  that "fits closely" the data? Some possibilities are (Eq. 9)

linear:  $g(x) = a_1 + a_2 x$ ,

quadratic:  $g(x) = a_1 + a_2 x + a_3 x^2$ ,

cubic:  $g(x) = a_1 + a_2 x + a_3 x^2 + a_4 x^3$ .



However, the choice of the approximating function  $g(x)$  to be used must depend on some knowledge of the behavior of the phenomenon under observation. For the oscilloscope trace in Figure 3, the behavior is clearly periodic and a function like (Eq. 10)

$$g(x) = a_1 \sin(x) + a_2 \cos(x).$$

would be a reasonable place to begin. Let us assume that the five points  $(0, -0.9)$ ,  $(\pi/4, 1.5)$ ,  $(\pi/2, 3.1)$ ,  $(3\pi/4, 3.0)$ , and  $(\pi, 1.1)$  are on the graph in Figure 3, so that the total error is


$$E = [a_1 \sin(0) + a_2 \cos(0) + 0.9]^2 + [a_1 \sin(\pi/4) + a_2 \cos(\pi/4) - 1.5]^2 + [a_1 \sin(\pi/2) + a_2 \cos(\pi/2) - 3.1]^2 + [a_1 \sin(3\pi/4) + a_2 \cos(3\pi/4) - 3.0]^2 + [a_1 \sin(\pi) + a_2 \cos(\pi) - 1.1]^2.$$

Next minimizing with respect to the two independent parameters  $a_1$  and  $a_2$  yields two algebraic expressions that can immediately be solved for  $a_1$  and  $a_2$  or

$$a_1 = 3.14 \text{ and } a_2 = -1.02,$$

so that (Eq. 11)

$$g(x) = 3.14 \sin(x) - 1.02 \cos(x),$$



which is shown in Figure 4 as the magenta curve.



UCES

# "Overdetermined Systems and Curve Fitting to Data"

**Course:** Methods and Analysis in UCES

**Chapter:** 1.0 Matrix Computations

**Section:** 1.8 Overdetermined Systems and Curve Fitting to Data

**Introduction:** overdetermined system

**Applied Area:** business forecasting, nonlinear heat diffusion

**Model:** least squares

**Method:** normal equations

**Implementation:** Maple, Matlab

**Assessment:** ill-conditioned normal equation

**Homework:**

**Prerequisites:**

1. **Math:** Algebra and matrices
2. **Computing:** Maple or Matlab
3. **Science:** Physics

**Objectives:**

1. **Math:** Minimization, normal equation
2. **Computing:** Overdetermined systems
3. **Science:** Nonlinear thermal properties

**General Information:** This is an introduction to curve fitting and to nonlinear aspects of heat transfer. The normal equations are introduced to solve the least squares problem, and the possibility of ill-conditioned normal equations is introduced.

**Contact Person:** R. E. White, NCSU, [white@math.ncsu.edu](mailto:white@math.ncsu.edu)

**Revision Date:** 7-28-94

**Copyright ©1994, R. E. White. All rights reserved.**

[Return to Table of Contents](#)



[Advanced Search](#) [Preferences](#) [Language Tools](#) [Search Tips](#)

number analysis and approximation least squares

Google Search

The "AND" operator is unnecessary -- we include all search terms by default.  
[[details](#)]

[Web](#) | [Images](#) | [Groups](#) | [Directory](#) | [News](#) |

Searched the web for **number analysis and approximation least squares**. Results 1 - 10 of about 154,000. See

### SINUM: Volume 34, Number 5

SIAM Journal on Numerical **Analysis**. Volume 34, **Number 5**. Contents. ...  
**Analysis and Approximation**

of Conservation Laws with Source Terms JM Greenberg, AY LeRoux, R ...  
epubs.siam.org/sam-bin/dbq/toc/SINUM/34/5 - 6k - Oct 21, 2003 - [Cached](#) -  
[Similar pages](#)

### Sponsored Links

#### Tile Numbers from Spain

High quality handmade products from Spain. Address Tiles, fans & more.  
[www.spanishplates.com](http://www.spanishplates.com)  
Interest: [Interest:](#)

### SINUM: Volume 39, Number 5

SIAM Journal on Numerical **Analysis**. Volume 39, **Number 5**. ... Finite  
Element **Approximation**

of a Degenerate Allen--Cahn/Cahn--Hilliard System John W. Barrett,  
James F ...

epubs.siam.org/sam-bin/dbq/toc/SINUM/39/5 - 5k - Oct 21, 2003 - [Cached](#) -  
[Similar pages](#)

[ [More results from epubs.siam.org](#) ]

#### Numerical Methods for

Least Squares Problems  
By Ake Bjorck. Buy new or used.  
Amazon.com  
Interest: [Interest:](#)

[See your message here...](#)

### Least Squares Method Curve Fitting Program to download. Nonlinear ...

... Nonlinear **Least Squares** Regression **Analysis** Back to the top. ... limited to the **number**  
of points ... **Least Squares** Fitting, **Least Squares Approximation** Installation and ...

Description: A nonlinear **least squares** fitting program that handles input errors or a covariance matrix on each...

Category: [Science](#) > [Math](#) > [Statistics](#) > [Software](#)

[www.prz.rzeszow.pl/~janand/](http://www.prz.rzeszow.pl/~janand/) - 21k - [Cached](#) - [Similar pages](#)

### [PDF] 6 Least Squares, Fourier Analysis, and Related Approximation ...

File Format: PDF/Adobe Acrobat - [View as HTML](#)

... One can imagine a **number** of situations where that ... relationship between **Least Squares**  
and Fourier **analysis** suggests a broad **approximation** algorithm involving ...

[astrwww.cwru.edu/personal/collins/numbk/chapt6.pdf](http://astrwww.cwru.edu/personal/collins/numbk/chapt6.pdf) - [Similar pages](#)

### Citations: and orthogonal least squares learning - Wang, Mendel, ...

... implies the setting of: the **number** of units ... we can apply the SingularSpectrum **Analysis**  
(SSA) 15 ... Fuzzy basis functions, universal **approximation**, and orthogonal ...

[citeseer.nj.nec.com/context/1102310/0](http://citeseer.nj.nec.com/context/1102310/0) - 16k - [Cached](#) - [Similar pages](#)

### Parsimonious Least Norm Approximation - Bradley, Mangasarian, ...

... 39] 35] This **number** can be bounded by the **number** of patterns N ... 1992 2: IEEE Transactions  
on Pattern **Analysis** and Machine ... Parsimonious **least norm approximation**. ...

[citeseer.nj.nec.com/220518.html](http://citeseer.nj.nec.com/220518.html) - 26k - [Cached](#) - [Similar pages](#)

[ [More results from citeseer.nj.nec.com](#) ]

### Numerical Analysis - Sample Programs

... 4.8 **Analysis** of Roundoff Error in the Gaussian Algorithm. ... 6.6 The B-Splines: Applications  
6.7 Taylor Series 6.8 Best **Approximation: Least-Squares** Theory 6.9 ...

[www.ma.utexas.edu/CNA/NA2/sample.html](http://www.ma.utexas.edu/CNA/NA2/sample.html) - 17k - Oct 21, 2003 - [Cached](#) - [Similar pages](#)

### [PDF] CONVERGENCE ANALYSIS OF THE LEAST-SQUARES ESTIMATES FOR INFINITE ...

File Format: PDF/Adobe Acrobat - [View as HTML](#)

... The method is based on the **analysis** of an ... to estimate only a finite **number** of coefficients  
of ... investigation of connection between the **approximation** by "long ...

Journal of the ACM -- 1961

... JH Wilkinson. Error **analysis** of direct methods of matrix inversion. ... Volume 8, **Number** 4, October 1961. ... Asymptotic behavior of the best polynomial **approximation**. ...  
[theory.lcs.mit.edu/~jacm/jacm61.html](http://theory.lcs.mit.edu/~jacm/jacm61.html) - 19k - [Cached](#) - [Similar pages](#)

Approximation via Subspaces: Analysis and Geometry

... **Approximation** via Subspaces: **Analysis** and Geometry. ... not even the given infinite **number** of basis ... obtained by the method of **least squares approximation** via finite ...  
[www.math.ohio-state.edu/~gerlach/math/BVtypset/node16.html](http://www.math.ohio-state.edu/~gerlach/math/BVtypset/node16.html) - 25k - Oct 21, 2003 - [Cached](#) - [Similar pages](#)

Google

Result Page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

number analysis and approximation least squares

Google Search

[Search within results](#)

Dissatisfied with your search results? [Help us improve.](#)

[Google Home](#) - [Advertise with Us](#) - [Business Solutions](#) - [Services & Tools](#) - [Jobs, Press, & Help](#)

©2003 Google

## Welcome to IEEE Xplore®

☐ Home☐ What Can I Access?☐ Log-outYour search matched **12336** of **978562** documents.A maximum of **500** results are displayed, **15** to a page, sorted by **Relevance** in **descending** order.

You may refine your search by editing the current search expression or entering a new one the text box.

Then click **Search Again**.

## Tables of Contents

☐ Journals & Magazines☐ Conference Proceedings☐ Standards

## Results:

Journal or Magazine = **JNL** Conference = **CNF** Standard = **STD**

## Search

☐ By Author☐ Basic☐ Advanced**1 Total least squares and constrained least squares applied to frequency domain system identification***Young, W.R.; Irwin, D.;*

System Theory, 1993. Proceedings SSST '93., Twenty-Fifth Southeastern Symposium on, 7-9 March 1993

Page(s): 285 -290

[\[Abstract\]](#) [\[PDF Full-Text \(388 KB\)\]](#) **IEEE CNF**

## Member Services

☐ Join IEEE☐ Establish IEEE Web Account☐ Access the IEEE Member Digital Library**2 Algorithmic engineering as a teaching aid for DSP***McWhirter, J.G.;*

Teaching of Digital Signal Processing, IEE Colloquium on, 16 Feb 1995

Page(s): 10/1 -10/3

[\[Abstract\]](#) [\[PDF Full-Text \(104 KB\)\]](#) **IEEE CNF****3 Robust estimation of 3-D motion parameters in presence of correspondence mismatches***Chaudhuri, S.; Chatterjee, S.;*

Signals, Systems and Computers, 1991. 1991 Conference Record of the Twenty Asilomar Conference on, 4-6 Nov. 1991

Page(s): 1195 -1199 vol.2

[\[Abstract\]](#) [\[PDF Full-Text \(376 KB\)\]](#) **IEEE CNF****4 A new bias-compensating least-squares method for identification of stochastic linear systems in presence of colored noise***Nguyen, H.L.; Sibille, P.; Garnier, H.;*

Decision and Control, 1993., Proceedings of the 32nd IEEE Conference on, 15-19 Dec 1993

Page(s): 2038 -2043 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(380 KB\)\]](#) **IEEE CNF**

---

**5 Fast least-squares curve fitting using quasi-rational splines**

*Flickner, M.; Hafner, J.; Rodriguez, E.J.; Sanz, J.L.C.;*

Image Processing, 1994. Proceedings. ICIP-94., IEEE International Conference ,  
Volume: 1 , 13-16 Nov. 1994

Page(s): 686 -690 vol.1

[\[Abstract\]](#) [\[PDF Full-Text \(464 KB\)\]](#) **IEEE CNF**

---

**6 Weighted least squares algorithm for continuous-time model**

*Gao, A.J.;*

Decision and Control, 1994., Proceedings of the 33rd IEEE Conference on ; Volu  
14-16 Dec. 1994

Page(s): 263 -264 vol.1

[\[Abstract\]](#) [\[PDF Full-Text \(124 KB\)\]](#) **IEEE CNF**

---

**7 A state-space approach to adaptive RLS filtering**

*Sayed, A.H.; Kailath, T.;*

Signal Processing Magazine, IEEE , Volume: 11 Issue: 3 , July 1994

Page(s): 18 -60

[\[Abstract\]](#) [\[PDF Full-Text \(3164 KB\)\]](#) **IEEE JNL**

---

**8 Behavior of the partial correlation coefficients of a least squares lattice filter in the presence of a nonstationary chirp input**

*Soni, T.; Zeidler, J.R.; Ku, W.H.;*

Signal Processing, IEEE Transactions on [see also Acoustics, Speech, and Signal Processing, IEEE Transactions on] , Volume: 43 Issue: 4 , April 1995

Page(s): 852 -863

[\[Abstract\]](#) [\[PDF Full-Text \(904 KB\)\]](#) **IEEE JNL**

---

**9 Optimal tight frames and quantum measurement**

*Eldar, Y.C.; Forney, G.D., Jr.;*

Information Theory, IEEE Transactions on , Volume: 48 Issue: 3 , March 2002

Page(s): 599 -610

[\[Abstract\]](#) [\[PDF Full-Text \(272 KB\)\]](#) **IEEE JNL**

---

**10 Stochastic  $H^\infty$  identification: an iteratively weighted least square algorithm**

*Rangan, S.; Ren, W.;*

Decision and Control, 1994., Proceedings of the 33rd IEEE Conference on , Volu

14-16 . 1994  
Page(s): 3374 -3379 vol.4

[\[Abstract\]](#) [\[PDF Full-Text \(396 KB\)\]](#) **IEEE CNF**

---

**11 Average power in digital CMOS circuits using least square estimation**  
*Murugavel, A.K.; Ranganathan, N.; Chandramouli, R.; Chavali, S.;*  
VLSI Design, 2001. Fourteenth International Conference on , 3-7 Jan. 2001  
Page(s): 215 -220

[\[Abstract\]](#) [\[PDF Full-Text \(452 KB\)\]](#) **IEEE CNF**

---

**12 Multichannel adaptive least squares-relating the 'Kalman' recursive squares (RLS) and least squares lattice (LSL) adaptive algorithms**  
*Lewis, P.S.;*  
Acoustics, Speech, and Signal Processing, 1988. ICASSP-88., 1988 International Conference on , 11-14 April 1988  
Page(s): 1926 -1929 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(292 KB\)\]](#) **IEEE CNF**

---

**13 Normalised order statistic LMS adaptive filtering**  
*Chambers, J.A.;*  
Non-Linear Filters, IEE Colloquium on , 18 May 1994  
Page(s): 3/1 -3/8

[\[Abstract\]](#) [\[PDF Full-Text \(240 KB\)\]](#) **IEEE CNF**

---

**14 Contraction mapping: an important property in adaptive filters**  
*Rupp, M.;*  
Digital Signal Processing Workshop, 1994., 1994 Sixth IEEE , 2-5 Oct. 1994  
Page(s): 273 -276

[\[Abstract\]](#) [\[PDF Full-Text \(300 KB\)\]](#) **IEEE CNF**

---

**15 Least square filters for approximating perfect reconstruction filter banks**  
*Spurbeck, M.S.; Scharf, L.L.;*  
Signals, Systems and Computers, 1994. 1994 Conference Record of the Twenty Asilomar Conference on , Volume: 2 , 31 Oct.-2 Nov. 1994  
Page(s): 996 -1000 vol.2

[\[Abstract\]](#) [\[PDF Full-Text \(384 KB\)\]](#) **IEEE CNF**

---

[1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [11](#) [12](#) [13](#) [14](#) [15](#) [16](#) [17](#) [18](#) [19](#) [20](#) [21](#) [22](#) [23](#) [24](#) [25](#) [26](#) [27](#) [28](#) [29](#)  
[33](#) [34](#) [\[Next\]](#)



**Welcome to IEEE Xplore®**

- ☐ Home
- ☐ What Can I Access?
- ☐ Log-out


**Tables of Contents**

- ☐ Journals & Magazines
- ☐ Conference Proceedings
- ☐ Standards

**Search**

- ☐ By Author
- ☐ Basic
- ☐ Advanced

**Member Services**

- ☐ Join IEEE
  - ☐ Establish IEEE Web Account
  - ☐ Access the IEEE Member Digital Library
-  [Print Format](#)

Your search matched **657** of **978562** documents.

A maximum of **500** results are displayed, **25** to a page, sorted by **Relevance** in **descending** order. You may refine your search by editing the current search expression or entering a new one the text b Then click **Search Again**.

numerical <and> analysis <and> least <and> squares

[Search Again](#)

**Results:**

Journal or Magazine = **JNL** Conference = **CNF** Standard = **STD**

---

**1 Novel numerical algorithm for overhead lines protection and adaptive autoreclosure**

*Terzija, V.V.; Radojevic, Z.M.; Koglin, H.-J.;*

Developments in Power System Protection, 2001, Seventh International Confer (IEE) , 9-12 April 2001

Page(s): 387 -390

[\[Abstract\]](#) [\[PDF Full-Text \(268 KB\)\]](#) **IEE CNF**

---

**2 The importance of initial-phase selection agreement of PN sequences the standpoint of numerical analysis and simulation of a DS-CDMA syste**

*Karkkainen, K.H.A.; Leppanen, P.A.;*

Global Telecommunications Conference, 1998. GLOBECOM 98. The Bridge to GI Integration. IEEE , Volume: 6 , 8-12 Nov. 1998

Page(s): 3295 -3301 vol.6

[\[Abstract\]](#) [\[PDF Full-Text \(464 KB\)\]](#) **IEEE CNF**

---

**3 The equivalence of TLS and correspondence analysis**

*Yao, K.; Lorenzelli, F.; Kong, J.;*

Acoustics, Speech, and Signal Processing, 1992. ICASSP-92., 1992 IEEE Intern Conference on , Volume: 5 , 23-26 March 1992

Page(s): 305 -308 vol.5

[\[Abstract\]](#) [\[PDF Full-Text \(328 KB\)\]](#) **IEEE CNF**

---

**4 Fixed and floating point error analysis of QRD-RLS and STAR-RLS adaptive filters**

*Raghunath, K.J.; Parhi, K.K.;*

Acoustics, Speech, and Signal Processing, 1994. ICASSP-94., 1994 IEEE Intern Conference on , Volume: iii , 19-22 April 1994



[\[Abstract\]](#) [\[PDF Full-Text \(256 KB\)\]](#) [IEEE CNF](#)

---

**5 A hybrid numerical method for orbit correction**

*White, G.; Himel, T.; Shoaee, H.;*

Particle Accelerator Conference, 1997. Proceedings of the 1997 , Volume: 2 , 12 May 1997

Page(s): 2425 -2427 vol.2

[\[Abstract\]](#) [\[PDF Full-Text \(264 KB\)\]](#) [IEEE CNF](#)

---

**6 An iterative algorithm for time-variant filtering in the discrete Gabor transform domain**

*Xiang-Gen Xia; Shie Qiani;*

Acoustics, Speech, and Signal Processing, 1997. ICASSP-97., 1997 IEEE International Conference on , Volume: 3 , 21-24 April 1997

Page(s): 2121 -2124 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(344 KB\)\]](#) [IEEE CNF](#)

---

**7 A numerically stable fast Newton type adaptive filter based on order u fast least squares algorithm**

*Youhua Wang; Ikeda, K.; Nakayama, K.;*

Acoustics, Speech, and Signal Processing, 1998. ICASSP '98. Proceedings of the IEEE International Conference on , Volume: 3 , 12-15 May 1998

Page(s): 1713 -1716 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(300 KB\)\]](#) [IEEE CNF](#)

---

**8 Efficient distance protection and adaptive autoreclosure numerical algorithm**

*Terzija, V.V.; Wehrmann, S.; Radojevic, Z.; Koglin, H.-J.;*

Transmission and Distribution Conference and Exposition, 2001 IEEE/PES , Volume: 28 Oct.-2 Nov. 2001

Page(s): 669 -674 vol.2

[\[Abstract\]](#) [\[PDF Full-Text \(303 KB\)\]](#) [IEEE CNF](#)

---

**9 A numerically stable fast Newton-type adaptive filter based on order recursive least squares algorithm**

*Youhua Wang; Ikeda, K.; Nakayama, K.;*

Signal Processing, IEEE Transactions on [see also Acoustics, Speech, and Signal Processing, IEEE Transactions on] , Volume: 51 Issue: 9 , Sept. 2003

Page(s): 2357 -2368

[\[Abstract\]](#) [\[PDF Full-Text \(827 KB\)\]](#) [IEEE JNL](#)

---

10 **Fast adaptive RLS algorithms: a generalized inverse approach and an**  
*Qiao, S.;*

Signal Processing, IEEE Transactions on [see also Acoustics, Speech, and Signal Processing, IEEE Transactions on], Volume: 39 Issue: 6, June 1991

Page(s): 1455 -1459

[\[Abstract\]](#) [\[PDF Full-Text \(400 KB\)\]](#) **IEEE JNL**

---

11 **Exterior moment method analysis of conducting scatterers by using interior Green's function and the method of least square**

*Sawaya, K.; Yatabe, S.; Adachi, S.;*

Antennas and Propagation, IEEE Transactions on, Volume: 40 Issue: 5, May 1

Page(s): 563 -565

[\[Abstract\]](#) [\[PDF Full-Text \(276 KB\)\]](#) **IEEE JNL**

---

12 **MAP and regularized constrained total least-squares image restoration**

*Mesarovic, V.Z.; Galatsanos, N.P.;*

Image Processing, 1994. Proceedings. ICIP-94., IEEE International Conference  
Volume: 3, 13-16 Nov. 1994

Page(s): 177 -181 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(380 KB\)\]](#) **IEEE CNF**

---

13 **Digital analysis of rotated images**

*Alliney, S.;*

Pattern Analysis and Machine Intelligence, IEEE Transactions on, Volume: 15  
Part 1, May 1993

Page(s): 499 -504

[\[Abstract\]](#) [\[PDF Full-Text \(564 KB\)\]](#) **IEEE JNL**

---

14 **Acoustic echo cancellation with the fast affine projection**

*Reed, M.J.; Hawksford, M.O.J.;*

Audio and Music Technology: The Challenge of Creative DSP (Ref. No. 1998/47  
Colloquium on, 18 Nov. 1998

Page(s): 16/1 -16/8

[\[Abstract\]](#) [\[PDF Full-Text \(568 KB\)\]](#) **IEEE CNF**

---

15 **Instantaneous error analysis of the least mean square algorithm**

*Milner, S.H.; Cryan, R.A.; Senior, J.M.; South, C.R.;*

Mathematical Aspects of Digital Signal Processing, IEEE Colloquium on, 10 Feb 1

Page(s): 4/1 -4/5

[\[Abstract\]](#) [\[PDF Full-Text \(352 KB\)\]](#) **IEEE CNF**

---

16 **IEEE Colloquium on 'New Directions in Adaptive Signal Processing' (D N .039)**

New Directions in Adaptive Signal Processing, IEE Colloquium on , 16 Feb 1993

[\[Abstract\]](#) [\[PDF Full-Text \(12 KB\)\]](#) [IEEE CNF](#)

---

17 **The effect of numerical stabilisation on AR spectral estimation**

Moore, A.; McLaughlin, S.;

New Directions in Adaptive Signal Processing, IEE Colloquium on , 16 Feb 1993

Page(s): 3/1 -3/9

[\[Abstract\]](#) [\[PDF Full-Text \(340 KB\)\]](#) [IEEE CNF](#)

---

18 **Frequency-domain penalized least-squares beamformer design for detection of breast cancer via microwave imaging**

Davis, S.K.; Xu Li; Bond, E.J.; Hagness, S.C.; Van Veen, B.D.;

Sensor Array and Multichannel Signal Processing Workshop Proceedings, 2002 Aug. 2002

Page(s): 120 -124

[\[Abstract\]](#) [\[PDF Full-Text \(609 KB\)\]](#) [IEEE CNF](#)

---

19 **Free-space millimeter-wave measurement of the complex dielectric constant of radome materials using nonlinear least-squares analysis**

Rao, B.R.;

Antennas and Propagation Society International Symposium, 1989. AP-S. Dige 26-30 June 1989

Page(s): 1534 -1537 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(128 KB\)\]](#) [IEEE CNF](#)

---

20 **On the convergence of least-squares estimates**

Nassiri-Toussi, K.; Ren, W.;

Decision and Control, 1992., Proceedings of the 31st IEEE Conference on , 16-1 1992

Page(s): 2233 -2238 vol.2

[\[Abstract\]](#) [\[PDF Full-Text \(457 KB\)\]](#) [IEEE CNF](#)

---

21 **Spline-based LMS method with constraints for spectrophotometric d correction**

Ben Slima, M.; Morawski, R.Z.; Barwicz, A.;

Instrumentation and Measurement Technology Conference, 1992. IMTC '92., 9t , 12-14 May 1992

Page(s): 594 -597

[\[Abstract\]](#) [\[PDF Full-Text \(268 KB\)\]](#) [IEEE CNF](#)

**22 A fast approximate RLS algorithm***Chansarkar, M.M.; Desai, U.B.;*

TENCON '93. Proceedings. Computer, Communication, Control and Power Engineering. 1993 IEEE Region 10 Conference on Issue: 0, 19-21 Oct. 1993  
Page(s): 532 -536 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(296 KB\)\]](#) **IEEE CNF**

---

**23 A wideband multichannel array sensor with a frequency domain adaptive filter structure***Goldstein, J.S.;*

Acoustics, Speech, and Signal Processing, 1993. ICASSP-93., 1993 IEEE International Conference on, Volume: 4, 27-30 April 1993  
Page(s): 520 -523 vol.4

[\[Abstract\]](#) [\[PDF Full-Text \(216 KB\)\]](#) **IEEE CNF**

---

**24 Wavelet transform domain LMS algorithm***Hosur, S.; Tewfik, A.H.;*

Acoustics, Speech, and Signal Processing, 1993. ICASSP-93., 1993 IEEE International Conference on, Volume: 3, 27-30 April 1993  
Page(s): 508 -510 vol.3

[\[Abstract\]](#) [\[PDF Full-Text \(240 KB\)\]](#) **IEEE CNF**

---

**25 Performance analysis of time domain block LMS algorithm***Hsieh, L.S.L.; Wood, S.L.;*

Acoustics, Speech, and Signal Processing, 1993. ICASSP-93., 1993 IEEE International Conference on, Volume: 3, 27-30 April 1993  
Page(s): 535 -538 vol.3


[\[Abstract\]](#) [\[PDF Full-Text \(256 KB\)\]](#) **IEEE CNF**

---

[1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [11](#) [12](#) [13](#) [14](#) [15](#) [16](#) [17](#) [18](#) [19](#) [20](#) [\[Next\]](#)

[Home](#) | [Log-out](#) | [Journals](#) | [Conference Proceedings](#) | [Standards](#) | [Search by Author](#) | [Basic Search](#) | [Advanced Search](#)  
[Join IEEE](#) | [Web Account](#) | [New this week](#) | [OPAC Linking Information](#) | [Your Feedback](#) | [Technical Support](#) | [Email Alerting](#)  
[No Robots Please](#) | [Release Notes](#) | [IEEE Online Publications](#) | [Help](#) | [FAQ](#) | [Terms](#) | [Back to Top](#)

Copyright © 2003 IEEE — All rights reserved



[Advanced Search](#) [Preferences](#) [Language Tools](#) [Search Tips](#)

number analysis and approximation least squares in physics

Google Search

"in" is a very common word and was not included in your search. [\[details\]](#)  
The "AND" operator is unnecessary -- we include all search terms by default.  
[\[details\]](#)

[Web](#) [Images](#) [Groups](#) [Directory](#) [News](#)

Searched the web for **number analysis and approximation least squares in physics**. Results 1 - 10 of about 4

### PHYS 251 - Numerical Analysis - Approximating Functions

... values of a function  $f(x)$  at a finite **number** of discrete ... of representing  $f(x)$  by  $p(x)$  is called a trigonometric **approximation**, and; ... **Least Square Data Analysis**. ...  
[www.physics.gmu.edu/~amin/phys251/Topics/NumAnalysis/ Approximation/approximatingFcn.html](http://www.physics.gmu.edu/~amin/phys251/Topics/NumAnalysis/Approximation/approximatingFcn.html) - 5k - [Cached](#) - [Similar pages](#)

### PHYS 251 - Data Analysis Basics

**Data Analysis** Basics. ... In this technique, we are not forcing equality at a certain **number** of points ... But again, this will be done to some order of **approximation**. ...  
[www.physics.gmu.edu/~amin/phys251/Topics/ DataAnalVisual/dataAnalysisBasics.html](http://www.physics.gmu.edu/~amin/phys251/Topics/ DataAnalVisual/dataAnalysisBasics.html) - 6k - [Cached](#) - [Similar pages](#)

### Least Squares Method Curve Fitting Program to download. Nonlinear ...

... Nonlinear **Least Squares** Regression **Analysis** Back to the top. ... limited to the **number** of points ... **Least Squares** Fitting, **Least Squares Approximation** Installation and ...  
Description: A nonlinear **least squares** fitting program that handles input errors or a covariance matrix on each...  
Category: [Science](#) > [Math](#) > [Statistics](#) > [Software](#)  
[www.prz.rzeszow.pl/~janand/](http://www.prz.rzeszow.pl/~janand/) - 21k - [Cached](#) - [Similar pages](#)

### [PDF] Least-Squares Fitting of a Hyperplane

File Format: PDF/Adobe Acrobat - [View as HTML](#)  
... used for illustration by York (1966), this **approximation** is good ... where  $? = n - m$  is the **number** of degrees ... Data Reduction and Error **Analysis** for The Physical ...  
[www.dsm.fordham.edu/~moniot/hyperplane.pdf](http://www.dsm.fordham.edu/~moniot/hyperplane.pdf) - [Similar pages](#)

### UCES Methods and Analysis Chap. 1.8: Overdetermined Systems and ...

... Course: Methods and **Analysis** in UCES. ... This matrix might have a large condition **number**. ... The **least squares** linear **approximation** model requires that the data be in ...  
[www.krellinst.org/UCES/archive/classes/ CNA/dir1.8/uces1.8.html](http://www.krellinst.org/UCES/archive/classes/ CNA/dir1.8/uces1.8.html) - 16k - [Cached](#) - [Similar pages](#)

### 3.?????

... Eigenvalue problems, Interpolation and polynomial **approximation**, Collocation methods ... Vector Bundle Theory, Euler **number**, Hopf Degree ... **Analysis** on Manifolds. ...  
[math.postech.ac.kr/eng/graduate/courseintro.htm](http://math.postech.ac.kr/eng/graduate/courseintro.htm) - 56k - [Cached](#) - [Similar pages](#)

### [PDF] ANALYSIS OF THE CONDITIONING NUMBER OF THE PLANE WAVE ...

File Format: PDF/Adobe Acrobat  
... in this paper with a reduced **number** of the ... A. de La Bourdonnaye, High frequency **approximation** of integral ... Mathematical Modelling and Numerical **Analysis**, 28, pp ...  
[www.dur.ac.uk/omar.laghrouche/eccomas2000.pdf](http://www.dur.ac.uk/omar.laghrouche/eccomas2000.pdf) - [Similar pages](#)

### IPAM - Multiscale Geometry and Analysis in High Dimensions

... In the past decade, a **number** of independent ... MGA provides in mathematical **analysis**, a complete ... **least squares approximation** in **approximati n** theory, data ...  
[www.ipam.ucla.edu/programs/mga2004/](http://www.ipam.ucla.edu/programs/mga2004/) - 16k - Oct 21, 2003 - [Cached](#) - [Similar pages](#)

### faculty

... functions with a finite **number** of discontinuities ... and computation of radial-basis-function

Google Search: number analysis and approximation least squares in physics <http://www.google.com/search?q=number+a...physics&hl=en&lr=&ie=UTF-8&start=0&sa=N>  
**approximation.** ... Fluids and plasmas; Modeling and **analysis** of nonlinear  
[www.math.tau.ac.il/am/faculty.htm](http://www.math.tau.ac.il/am/faculty.htm) - 36k - [Cached](#) - [Similar pages](#)

**The Math Forum - Math Library - Numerical Analysis**

... routines, data structures, random **number** generators, and ... A paper on **approximation**  
and various resources ... applied mathematics, numerical **analysis**, fluid dynamics ...  
[mathforum.org/library/topics/num\\_analysis/](http://mathforum.org/library/topics/num_analysis/) - 36k - Oct 21, 2003 - [Cached](#) - [Similar pages](#)

Google

Result Page:    1   2   3   4   5   6   7   8   9   10    [Next](#)

[Search within results](#)

Dissatisfied with your search results? [Help us improve.](#)

[Google Home](#) - [Advertise with Us](#) - [Business Solutions](#) - [Services & Tools](#) - [Jobs, Press, & Help](#)

©2003 Google